Author Index

Allmendinger, H., see Koppenhoefer, B. 89
Armani, E.
——, Barazzoni, L., Dossena, A. and Marchelli, R. Bis(L-amino acid amidato)copper(II) complexes as chiral eluents in the enantiomeric separation of D,L-dansylamino acids by reversed-phase high-performance liquid chromatography 287
——, Dossena, A., Marchelli, R. and Virgili, R. Copper(II) complexes of diamino-diamido-type ligands as chiral eluents in the enantiomeric separation of D,L-dansylamino acids by reversed-phase high-performance liquid chromatography 275
Armstrong, D. W., see Han, S. M. 376
Barazzoni, L., see Armani, E. 287
Barkley, R. M., see DeAngelis, J. J. 125
Batten, C. F., see Wentworth, W. E. 45
Bellama, J. M., see Tierney, E. J. 229
Bhatia, A., see Jain, R. 454
Bhatnagar, D., see McCormick, S. P. 400
Blouin, F. A.
—— and Zarins, Z. M. High-performance liquid chromatography of cottonseed flavonoids 443
Bowers, E., see McCormick, S. P. 400
Brinckman, F. E., see Tierney, E. J. 229
Brooks, C. J. W.
—— and Cole, W. J. Cyclic di-*tert.*-butylsilylene derivatives of substituted salicylic acids and related compounds. A study by gas chromatography–mass spectrometry 13
Burford, R. P., see Haken, J. K. 207
Cagniant, E., see Tillier, C. 406
Cai, H., see Yamaguchi, J. 183
Chang, L. P., see Koppenhoefer, B. 89
Chen, E. C. M., see Wentworth, W. E. 45
Cheng, L. B., see Koppenhoefer, B. 89
Cohen, S. M., see Tibbels, T. S. 448
Cole, W. J., see Brooks, C. J. W. 13
Daussant, J., see Lecommandeur, D. 436
DeAngelis, J. J.
——, Barkley, R. M. and Sievers, R. E. Coupling liquid chromatography with redox chemiluminescence detection 125
Devaux, Ph., see Tillier, C. 406
Dossena, A., see Armani, E. 275, 287
Driscoll, J. N., see Duffy, M. 73
——, Duffy, M. and Pappas, S. Capillary gas chromatographic analysis with the far-UV absorbance detector 63
Duffy, M., see Driscoll, J. N. 63
——, Driscoll, J. N., Pappas, S. and Sanford, W. Analysis of ppb levels of organics in water by means of purge-and-trap, capillary gas chromatography and selective detectors 73
Egeler, E., see Jacob, K. 171
Eng, G., see Tierney, E. J. 229
Eriksson, A., see Qureshi, G. A. 197
Ewing, A. G., see Wallingford, R. A. 299
Foley, J. P. Separation of *m*- and *p*-cresol and other methylated phenols in shale oil by reversed-phase liquid chromatography with electrochemical detection 347
Fox, A., see Gilbart, J. 323
Galensa, R., see Krause, M. 417
Games, D. E., see Morgan, E. D. 165
Gilbart, J.
——, Harrison, J., Parks, C. and Fox, A. Analysis of the amino acid and sugar composition of streptococcal cell walls by gas chromatography–mass spectrometry 323
Guillemin, C. L. New concept in chromatographic instrumentation. The probe-process liquid chromatograph 1
Haken, J. K.
——, Harahap, N. and Burford, R. P. Gas chromatographic procedure for the qualitative analysis of silicone polyester resins after alkaline fusion 207
Han, S. M.
——, Han, Y. I. and Armstrong, D. W. Structural factors affecting chiral recognition and separation on β-cyclodextrin bonded phases 376
Han, Y. I., see Han, S. M. 376
Hanai, T., see Yamaguchi, J. 183
Harahap, N., see Haken, J. K. 207
Harrison, J , see Gilbart, J. 323
Hengartner, H., see Schott, H. 115
Herrmann, K., see Klick, S. 335
Hicks, K. B.
—— and Hotchkiss, Jr., A. T. High-performance liquid chromatography of plant-derived oligosaccharides on a new cation-exchange resin stationary phase: HPX-22H 382

Hill, E.
——, Humphreys, E. and Malcolme-Lawes, D. J. Electrochemiluminescence as a detection technique for reversed-phase high-performance liquid chromatography. IV. Detection of fluorescent derivatives 394
Hotchkiss, Jr., A. T., see Hicks, K. B. 382
Humphreys, E., see Hill, E. 394
Jacob, K.
——, Kossien, I., Egeler, E. and Knedel, M. Detection of non-typical porphyrin isomers in human urines by ion-pair reversed-phase high-performance liquid chromatography 171
Jain, R.
—— and Bhatia, A. Thin-layer chromatographic separation of some sulpha drugs using acetoacetanilide as a coupling agent 454
Johannesen, R. B., see Tierney, E. J. 229
Jones, K.
—— and Malcolme-Lawes, D. J. High-performance ultraviolet absorption detector for liquid chromatography. II. A prototype detector 387
Kaplan, L., see Metz, P. A. 31
Karmen, A., see Lam, S. 81
Klick, S.
—— and Herrmann, K. Analysis of naturally occurring hydroxybenzoic acid glucosides and hydroxybenzoyl glucoses and their behaviour on a polyamide column 335
Knedel, M., see Jacob, K. 171
Koppenhoefer, B.
——, Allmendinger, H., Chang, L. P. and Cheng, L. B. Resolution of stereoisomers of dipeptides by gas chromatography on Chirasil-Val 89
Kossien, I., see Jacob, K. 171
Krause, M.
—— and Galensa, R. Direct enantiomeric separation of racemic flavanones by high-performance liquid chromatography using cellulose triacetate as a chiral stationary phase 417
Krob, E. J., see Walter, H. 261
Lam, S.
——, Malikin, G. and Karmen, A. High-performance liquid chromatography of hydroxysteroids detected with post-column immobilized enzyme reactors 81
Lecommandeur, D.
——, MacGregor, A. W. and Daussant, J. Purification of germinated barley α-amylase isozymes and limit-dextrinase by chromatofocusing and affinity chromatography 436
Leitner, B., see Schott, H. 115
Liesch, J. M., see Metz, P. A. 31
Limero, T., see Wentworth, W. E. 45
Lipinski, J.
—— and Stan, H.-J. CAPA —Computer aided pesticide analyses. Computer program for the automated evaluation of chromatographic data for residue analysis foods 213
Liu, G.
Oxidation of thiols within gas chromatographic columns 372
McCormick, S. P.
——, Bowers, E. and Bhatnagar, D. High-performance liquid chromatographic procedure for determining the profiles of aflatoxin precursors in wildtype and mutant strains of *Aspergillus parasiticus* 400
McCune, J. E., see Pirkle, W. H. 311
MacGregor, A. W., see Lecommandeur, D. 436
Malcolme-Lawes, D. J., see Hill, E. 394
——, see Jones, K. 387
Malikin, G., see Lam, S. 81
Marchelli, R., see Armani, E. 275, 287
Mauri, P., see Pietta, P. 431
Metz, P. A.
——, Omstead, D. R., Kaplan, L., Liesch, J. M., Stearns, R. A. and Vandenheuvel, W. J. A. Characterization of a lipid-rich fraction synthesized by *Streptomyces avermitilis* 31
Morgan, E. D.
——, Murphy, S. J., Games, D. E. and Mylchreest, I. C. Analysis of ecdysteroids by supercritical-fluid chromatography 165
Murphy, S. J., see Morgan, E. D. 165
Mylchreest, I. C., see Morgan, E. D. 165
Omstead, D. R., see Metz, P. A. 31
Pace, M., see Pietta, P. 431
Pappas, S., see Driscoll, J. N. 63
——, see Duffy, M. 73
Parks, C., see Gilbart, J. 323
Pietta, P.
——, Mauri, P. and Pace, M. High-performance liquid chromatographic method for the determination of the esterase activity of subtilisin and kallikrein 431
Pirkle, W. H.
—— and McCune, J. E. An improved chiral stationary phase for the facile separation of enantiomers 311
Potschka, M.
Size-exclusion chromatography of polyelectrolytes. Experimental evidence for a general mechanism 239

Qureshi, G. A.
—— and Eriksson, A.
Determination of clenbuterol and mabuterol in equine plasma by ion-pair liquid chromatography with electrochemical detection. Chromatographic and electrochemical characteristics 197
Roeraade, J.
Factors affecting sample transfer from microlitre syringes 367
Ruane, R. J.
—— and Wilson, I. D.
Ion-pair reversed-phase thin-layer chromatography of basic drugs using sulphonic acids 355
Sanford, W., see Duffy, M. 73
Schott, H.
——, Leitner, B., Schwendener, R. A. and Hengartner, H.
Chromatography of functionalized liposomes and their components 115
Schurig, V.
Enantiomer analysis by complexation gas chromatography. Scope, merits and limitations 135
—— and Nowotny, H.-P.
Separation of enantiomers on diluted permethylated β-cyclodextrin by high-resolution gas chromatography 155
Schwendener, R. A., see Schott, H. 115
Segura, R.
Preparation of fatty acid methyl esters by direct transesterification of lipids with aluminium chloride–methanol 99
Sievers, R. E., see DeAngelis, J. J. 125
Smith, R. A., see Tibbels, T. S. 448
Stan, H.-J., see Lipinski, J. 213
Stearns, R. A., see Metz, P. A. 31
Tibbels, T. S.
——, Smith, R. A. and Cohen, S. M.
Determination of saccharin in diet and biological materials 448
Tierney, E. J.
——, Bellama, J. M., Eng, G., Brinckman, F. E. and Johannesen, R. B.
Total molecular surface areas as a predictor for reversed-phase high-performance liquid chromatography in various organotin systems. Importance and applications of carbon hybridization summaries 229
Tillier, C.
——, Cagniant, E. and Devaux, Ph.
Determination of halofuginone in poultry feeds by high-performance liquid chromatography 406
Vaara, M.
Analytical and preparative high-performance liquid chromatography of the papain-cleaved derivative of polymyxin B 423
Vandenheuvel, W. J. A., see Metz, P. A. 31
Virgili, R., see Armani, E. 275
Wallingford, R. A.
—— and Ewing, A. G.
Retention of ionic and non-ionic catechols in capillary zone electrophoresis with micellar solutions 299
Walter, H.
—— and Krob, E. J.
Partitioning behavior of erythrocytes in aqueous two-phase systems containing hydroxypropyl starch and polyethylene glycol 261
Weinstein, L. H., see Wettlaufer, S. H. 361
Wentworth, W. E.
——, Limero, T., Batten, C. F. and Chen, E. C. M.
Non-radioactive electron-capture detector operating in the pulsed mode 45
Wettlaufer, S. H.
—— and Weinstein, L. H.
Quantitation of polyamines using thin-layer chromatography and image analysis 361
Wilson, I. D., see Ruane, R. J. 355
Yamaguchi, J.
——, Hanai, T. and Cai, H.
Selectivity related to carbon loading and end-capping of octadecyl-bonded silica gels in reversed-phase liquid chromatography 183
Zarins, Z. M., see Blouin, F. A. 443